



RESEARCH ARTICLE

NON-EXTRACTIVE SPECTROPHOTOMETRIC DETERMINATION OF NI(II) IN PLANT SAMPLES USING SALICYLALDEHYDE ISONICOTINOYLHYDRAZONE

Renuka M^{1*} and Hussain Reddy K²

¹Chemistry, Loyola Degree Collage, Y.S.R (Dist)

²Department, of Chemistry S.K.University, Anantapur (Dist)

ARTICLE INFO

Article History:

Received 19th, December, 2015

Received in revised form 28th,
December, 2015

Accepted 13th, January, 2015

Published online 28th,

January, 2016

Key words

Spectrophotometric determination,
Salicylaldehyde isonicotinoylhydrazone,
plant Samples, molar absorptivity
and sandell's sensitivity

ABSTRACT

A very simple, highly selective and non-extractive spectrophotometric determination for the trace amounts of nickel (II) has been developed. Salicylaldehyde isonicotinoylhydrazone (SAINH) has been proposed as a new analytical reagent for the direct non- extractive spectrophotometric determination of nickel (II). The reagent reacts with nickel in an basic medium (pH 7.5, sodium acetate and acetic acid) to form a yellow coloured 1:1(M:L) Complex. The reaction is instantaneous and the maximum absorption was obtained at 385 nm and remains stable for 1h. The molar absorptivity and sandell's sensitivity were found to be $1.81 \times 10^4 \text{ L mol}^{-1} \text{ cm}^{-1}$ and $0.32 \mu\text{g cm}^{-2}$ respectively. Linear calibration graphs were obtained for 1.0 -10.0 $\mu\text{g/ml}$ of nickel (II). The method is highly selective for nickel and successfully used for the determination of nickel in several plant samples.

© Copy Right, Research Alert, 2016, Academic Journals. All rights reserved.

INTRODUCTION

Nickel is a silvery white, lustrous metal with a slight golden tinge. It is hard and ductile. Pure nickel shows a significant chemical activity, can be observed when it is powdered to maximize the exposed surface area on which reactions can occur, but larger pieces of the metal are slow to react with air at ambient conditions due to the formation of a protective oxide surface.

Naturally occurring nickel is composed of 5 stable isotopes ^{58}Ni , ^{60}Ni , ^{61}Ni , ^{62}Ni and ^{64}Ni and ^{58}Ni is the most abundant (68.08%), ^{62}Ni is the most stable nuclide of all the existing elements [1]. The common oxidation states of nickel is +2, but compounds of Ni, Ni^+ and Ni^{3+} are well known and Ni^{4+} has been demonstrated [2]. Because of the nickel slow rate of oxidation at room temperature, it is considered as a corrosion resistant. Because of this it has been occasionally used as substitute for decorative silver. It is used for plating metals such as iron and brass, for chemical apparatus and in certain alloys that retain a high silvery polish such as German silver. As a compound nickel used as catalyst for hydrogenation. Enzymes of some microorganisms and plants contain nickel as an active site which makes the metal an essential nutrient for them. It is used in many specific and recognizable industrial and consumer products including stainless steel, alnico magnets, coinage, rechargeable batteries, electric guitar strings, microphone capsules and special and alloys. Nickel is absorbed by inhalation and ingestion.

The immediate effects of nickel carbonyl exposure are respiratory tract irritation and neurological effects, nausea, vomiting, diarrhea and head ache. Delayed effects include pulmonary oedema, pneumonia and in severe cases death.

In natural water environment, nickel concentration is generally small because of its occurrence in a colloidal form. It is easily accumulated in the biological environment, particularly in the phytoplankton or other aquatic plants which are sensitive bio indicators of water pollution. Since nickel is extensively used in the preparation of alloys and catalysts, its determination may be considered as interesting research activity. Various spectrophotometric used for the determination of nickel are summarized in Table 1. [3-13]

MATERIALS AND METHODS

Preparation of Sainh

Salicylaldehyde (1.05 ml, 0.01 mol) and isoniazid (0.69g, 0.01 mol dissolved in 3 ml of ethanol) were mixed in a clean round bottom flask. Suitable quantity of (~10ml) of ethanol was added to the reaction mixture and refluxed with stirring for 2 hrs. A pale yellow coloured product was separated out on cooling. It was collected by filtration, washed several times with hot water. This compound was recrystallized from methanol and dried in vacuo. Yield 2.8 gm, melting point is 250 – 252°C.

Table 1 Spectrophotometric methods for the determination of nickel (II) with different Reagents

Name of the reagent	max (nm)	pH	Beers law(ppm)	Molar absorptivity	M:L	Ref
				() L mol ⁻¹ cm ⁻¹		
Molybdophosphoric acid	490	4.3	0.0-5.0	0.13x10 ⁴	1:01	3
4-(5-utiro-2-thiozolyazu resorcinol)	540	6.5	4	1.0x10 ⁴	1:01	4
Sodium isoamylxanthate	360	6.0-8.0	0.2-2.7	1.2x10 ⁴	1:02	5
Pthalaldehyde dithiosemicarbazone (PADT)	400	7.5-9.0	Upto4.5	0.81x10 ⁴	1:01	6
Quinolaldehyde thiosemicarbazone (QAH)	460	7.5	5.0-25.0	1.6x10 ⁴	1:02	7
Salicylaldehyde thiosemicarbazone (SAT)	370	6.5-7.0	0.5-5.0	0.98x10 ⁴	1:01	8
Acenaphthaquinone thiosemicarbazone (ANQT)	420	6.3-8.4	Upto14.7	0.28x10 ⁴	1:02	9
Diacetylmonoxime-4-phenyl-3-thiosemicarbazone (DAMOPT)	375	5.2-10.0	-	1.7x10 ⁴	-	10
2-hydroxy-4-isopropoxy acetophenone thiosemicarbazone (HIAT)	400	9	1.6-4.4	0.08x10 ⁴	1:02	11
2-(2-Furanylmethylene) hydrazine carbothioamide (FMHC)	440	5.8	0.36-6.0	0.72x10 ⁴	1:02	12
1-phenyl-1,2-propanedione-2-thiosemicarbazone (PPDOT)	395	3.0-6.0	1.8-3.7	1.2x10 ⁴	1:02	13
Salicylaldehyde isonicotinoylhydrazone (SAINH)	378	1.0-10.0	7.5	1.81x10 ⁴	1:01	Present method

Preparation of SAINH solution

It was prepared by taking 0.060 gm of SAINH reagent substance in a 25–ml standard flask. The reagent substance was dissolved in 10 ml of DMF and diluted up to the mark with the same solvent.

Preparation of Nickel (Ii) Ion Solution

Stock solution (1 × 10⁻²M) of Ni(II) was prepared by dissolving requisite quantity (0.25 g) of nickel sulphate (NiSO₄.7H₂O) in doubly distilled water containing a few drops of conc.H₂SO₄ and made upto 100-ml volumetric flask. The stock solution was standardized gravimetrically[14].

Recommended Procedure

An aliquot of the solution containing nickel in optimum concentration range , 10 ml of buffer solution (pH 7.5) and 1 ml of 0.01M reagent solution were combined in 25 – ml volumetric standard flask and resulting solution was diluted to the mark with distilled water. The absorbance of the solution was measured at 385 nm against reagent (SAINH) blank. The measured absorbance was used to compute the amount of nickel from predetermined calibration plot.

Preparation of Plant Samples

Freshly collected samples were cleaned and dried for one hour in the open air protecting from mineral contamination. The drug samples were finally powdered in a mortar. The powdered material was brought into solution by wet ashing method.

Instrumentation

Perkin – Elmer (Lamda 25), UV – Visible spectrophotometer equipped with 1.0 cm (path lenth) quartz cell and Elico model LI- 610 pH meter were used in the present study

RESULTS AND DISCUSSIONS

The reagent (SAINH) are easily obtained by condensation reaction and is shown in Fig 1.




Figure 1 Condensation reaction of SAINH

It has been characterized using IR, NMR and Mass spectral data. The infrared spectrum of SAINH showed bands (cm⁻¹) at 3271(m), 3120(m), 3051(m), 1649(s), 1619(s), 1535(m), 1473(s), 1290(s), 768(s), 711(s) and 682(m), are respectively assigned to (NH) secondary stretching, (C-H) stretching, (isoniazid), (C-H) aromatic stretching (salicylaldehyde), (C=O) isonicotinoyl Stretching, (N-H) plane bending, (C- C) aromatic ring stretching, (C-H) assymmetric bending, (C-N) streching vibrations, (C-H)

aromatic oop bend (salicylaldehyde), and (N-N) stretch and (C-H) Aromatic oop bending respectively. This spectrum is given in Fig 2.




Figure 2 Infrared Spectrum of SAINH in KBr disc

¹H – NMR spectra (DMSO – d₆) showed signals (ppm) at 6.65, 7.786-7.801 and 8.33 due to - CH₃, - OH (aliphatic), phenolic and - NH₂ protons of SAINH respectively. NMR spectrum is given in Fig 3.




Figure 3 ¹H-NMR Spectrum of SAINH in DMSO – d₆ medium

Massspectra shows molecular ion signal at m/z 241. Other peaks are observed at m/z values of 240 and 163 respectively due to the loss of -H and -C₅H₄N radicals. Based on above spectral data the structure of the reagent is given in Fig 4.




Figure 4 Mass Spectrum of SAINH in DMSO – d₆ medium

Absorption spectra of 2x10⁻⁵ solution of SAINH at different pH values were recorded and pKa values were determined spectrophotometrically using Phillip and Merrit method [15]. The bathochromic shift from 295 – 365 nm indicates that in solution on increasing pH the >C=O group of the reagent (SAINH) is enolised and dissociated. The values of SAINH are 4.2(pK₁) and 9.0

(pK₂) respectively. The pK₁ and pK₂ values are presumably due to keto – enol tautomerism and deprotonation of - NH group respectively.

A 0.01M solution of reagent is stable for 1h. The colour reactions of reagent with nickel (II) are instantaneous at room temperature. The order of addition of nickel ion, reagent and buffer has no effect on the absorbance of the complex. Various physico – chemical and analytical characteristics of nickel complex are summarized in Table 1. Stoichiometry of the complex (M : L = 1 : 1) was determined by Job’s continuous variation method and molar ratio method. Sodium acetate (1M) –acetic acid (0.1M) buffer (pH 7.5) is used in these studies. The dissociation constant () and concentration (c) of the reagent at intersecting point were used in the calculation of stability constant of the complex. Stability constant of the complex 1: 1 (M : L) complex is given by 1- / ²c. The predicted structure of Ni (II) – SAINH is given in Fig 5.




Figure 5 Predicted structure of Ni – SAINH complex

Tolerance Limits of Foreign Ions

The effect of foreign ions which often accompany nickel has been studied by adding different amounts of foreign ions to fixed amount of nickel (II) of ±2% in the absorbance of the reaction mixture was considered tolerable. The results are given in Table 2. Larger amounts of Cd(II) and Co(II) do not interfere in the presence of masking agents. Interference of Cd(II) and Co(II) are masked with thiocyanate.

Table 2 Tolerance limit of foreign ions in the determination of 4.69 µg/ml of Nickel

Ion added	Tolerance limit µg/ml	Ion added	Tolerance limit µg/ml
Citrate	544.2	U(VI)	95.2
Iodide	504.2	W(VI)	59
Phosphate	392	Cd(II)	27a
Oxalate	352	Mn(II)	22
Tartrate	296	Cr(VI)	12.5
Nitrate	248	Pb(II)	8.2
Sulphate	192	Co(II)	2.4a
Bromide	160	As(I)	4.1
Chloride	142	Ti(II)	2.4
Carbonate	120	Pt(IV)	2.0
Acetate	94	Pd(II)	0.90
Fluoride	78	Zn(II)	0.23
Thiourea	31		

a: Masked with 200µg/ml of thiocyanate

Applications

The present method was successfully applied for the determination of nickel in various plant samples and the results were presented in Table 3.

Table 3 Amount of nickel (%) present in various plant samples

Sample	Amount of Nickel*		Error
	in µg/ml		
	Taken	Found	
Eucalyptus leaves	1.063	1.060	0.28%
Pisium Sativum	2.051	2.062	0.53%
Mangifera indica leaves	2.226	2.206	0.86%
Azadirachta indica leaves	1.527	1.520	0.45%

* Average of three determinations

CONCLUSION

The synthesized reagent salicylaldehyde isonicotinoyl hydrazone (SAINH) is characterized by analytical and spectral studies. The reagent forms a yellow coloured complex with nickel (II). The Ni (II)-SAINH Complex structure is predicted and various physico-chemical and analytical characteristics are determined. This reagent SAINH is successfully used for the determination of nickel (II) in various plant samples.

Acknowledgement

The authors thank M.Subbalakshmi of IICT, Hyderabad for her help in recording IR, Mass and NMR spectra of reagent samples.

References

1. M.P.Fewell, The atomic nuclide with the highest mean binding energy *American journal of physics*. 2011, 63(7), 653-58.

2. Green Wood, N.Norman, Earn Shaw, Alan. Chemistry of the elements 1997, (2nd Ed.)
3. R.L. Heller, J.C.Guyon., *Talanta*, 1970, 17(9), 865-71.
4. A.K. Alikhodzheer, D.N. Pachadzha, M.Yusuprov, Z.Yu, Ana. Khim, 1985, 40(1), 2021.
5. A.K. Malik, W. Fawbel, B.S. Sedal and A.L.J. Rao., *Chem. Environ. Res.* (Review), 2000, in press.
6. H. Sanke Gouda, K.N. Thimmaiah and S. Maqbool Ahmed., *India.j.chem*, 1982, 2, 327.
7. D.V. Khasnis and V.M. Shinde., *J.indian. chem.soc*. 1984, 61, 275.
8. M. Subbarami Reddy, Ph.D. Thesis, S.K. University, Anantapur, India, 1982.
9. S. K. Singh, M. Kamini and S. K. Sindhvani., *J. Chinese. Chem.,soc.*, 1982, 29, 131
10. J.M. Cano Pavon, A.Garcia de Torres and F. Pino., *An. Quim.* 1978, 14, 24.
11. V.K, Desai, K.K. Desai., *Orient. J. chem...* 1996, 12, 203.
12. F. Salinas, J.C> Jiminez Sanchez and J.L. Lemus., *Bull.Soc.Chim. Beig*, 1986, 95,293.
13. H. Sanke Gouda, K.N. Thimmaiah and S. Maqbool Ahmed., *India.j.chem*. 1983,22,551.
14. A.I. Vogel., *A text book of quantitative inorganic Analysis*. 3rd edn, ELBS and Longman, 1975, p.325.
15. J.P. Phillips and L.L. Merrit., *J. Ame. Chim.Soc.* 1978, 70,410.
