



PERCENTAGE TO CONTROL FUNGAL DISEASES OF BRINJAL BY USING DIFFERENT LEAF EXTRACT

Anarse P. S.,

Dept. of Botany Gandhi College, Kada, Tq. Ashti Dist. Beed(M S)

ARTICLE INFO

Article History:

Received 15th July, 2020

Received in revised form 7th

August, 2020

Accepted 13th September, 2020

Published online 28th October, 2020

Key words:

pathogen, fungal diseases, leafextract, fungicides etc.

ABSTRACT

Now a days we know that due to the environmental condition atmosphere is not favorable for any crops, so that for healthy and more production we are spraying different fungicides, insecticides and etc. Spraying things are not good for human beings for that we are using different leaf extracts, This method is very easy and economically beneficially to producer. The Brinjal consist of different fungal diseases to control we are using different leaf extracts which affects to the plants growth and yield to a best of genetic potential. In present study , the pathogenic fungus was isolated from infected plant parts and identified based on morphological and cultural characters as *Fusarium Solanif.sp, melongenae* .Hossain KS and Bashar MA(2011).The different plant extracts viz, *Azardichataindica, Argemone mexicana, Datura stramonium, Ipomoea fistulosawere* tested to control brinjal pathogen. Different concentrations 5, 10, 15 and 20% of plant extracts was used in the study .Alemu Nega (2014). Among the different extracts 20% of *Azardiachtaindica* was found most effective followed by *Argemone mexicana, Datura stramonium, Ipomoea fistulosa*.

Copyright©2020 Anarse P. S. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Brinjal or eggplant (*Solanum melongena*L.) is an important crop of sub-tropics and tropics. The name brinjal is popular in Indian subcontinents and is derived from Arabic and Sanskrit whereas the name eggplant has been derived from the shape of the fruit of some varieties, which are white and resemble in shape to chicken eggs. It is also called aubergine (French word) in Europe. Jagannathan R. and Narasimhan (1988).The egg plant is native of India .Brinjal is grown as an important vegetable crop in all over world. It is grown in India over an area of 0.4 million hectares with an annual production of 7.8 million tonnesHossain KS and Bashar MA (2011). Among the different diseases that attack brinjal crop , wilt has become a major disease causing significant reduction in yield. The wilt of brinjal is characterized by yellowing of foliage drooping of apical shoot to ultimate death of whole plant. The pathogen is a soil inhabiting fungus and forms in the senescing tissues of the diseased plant and may survive in the soil for many years . There are many methods which are presently being used to control various plant pathogens including wilt pathogen such as physical, chemical, biological, cultural etc. Effective and efficient management of crop disease is generally achieved by the use of synthetic pesticides [2].due to increased awareness about the risks involved in use of pesticides, much attention is being focused on the alternative methods of pathogen control . The spiraling up cost chemical fungicides particularly in those countries where pesticide are imported. Pollution to soil,

water, air by the accumulation of obnoxious chemicals residues due to continuous use of fungicides and development of resistance races to these chemicals are therefore now facing the scientist to look for methods which are ecologically , friendly, safe and specific for pathogens . The recurrent and indiscriminate use of fungicides have posed a serious threat to human health and to the existing human eco geographical conditions as some of them have already been proved to be either mutagenic , carcinogenic or teratogenic. Keeping in view the drawback of chemical management of plant disease, the use of plant extracts in the management of plant disease is gaining importance.

MATERIALS AND METHODS

Roots and plant parts were collected from infected brinjal parts showing characteristic symptoms of pathogens from the field . Plant parts were sampled from the late winter -early autumn. The plant parts were then examined under microscope to confirm the presence of respective pathogen *Fusarium solani f. sp. Melongenae* and the infected plant parts were cut into pieces (2-3 mm), surface sterilized with 0.1% mercuric chloride solution for 30 seconds. The isolation was made from root as well as from the foliar parts of wilted Brinjal plants. The plant parts were washed three times with sterilized distilled water and then were transferred aseptically on Potato Dextrose Agar (PDA) media. After the development of the fungal colonies stock cultures were prepared using PDA in test

*Corresponding author: **Anarse P. S.,**

Dept. of Botany Gandhi College, Kada, Tq. Ashti Dist. Beed(M S)

tubes and stored in refrigerator at 4°C .Brinjal pathogen was isolated from infected brinjal plants and was identified.

Plants used in the present study are *Azardiachtaindica* which belongs to the family *Meliaceae* commonly knows as “ neem”. The plant is found throughout India and its derivatives are of great use in agriculture, public health, medicines, cosmetics and many more . The leaves, bark, seed and flowers are bitter , astringent, acid, depurative, refrigrant, demulcent, insecticidal, expectorant liver tonic, etc. An important of application of neem products in agriculture is their ability to nitrogen release from the nitrogenous fertilizers.

Ocimumsanctum, commonly known as “tulsi” belongs to the family *Lamiaceae* found throughout India. The plant is much erect, branched, softly pubescent under shrub, 30-60 cm high with red or purple sub-quadrangular branches, leaves simple, opposite , elliptic, whole plants is used as medicine for various diseases. *Tulsi* leaves contain a bright yellow volatile oil which is reported to possess antibacterial properties and acts as insecticide.

Datura stramonium plant is said to have been used as a narcotic as early as A.D. 37. It is one of the favourite source of “knockout drops” . It contain on alkaloid scopolamine which is said to produce hallucinatory effects. The smoking of the narcotic produces hallucinatory effects. The smoking of the narcotic produces pronounced diversions in ideas, emotions and even perception . The leaves and tops of *stramonium* are mixed with lobelia herb, lobelia inflata , often called Indian tobacco , to make asthma powders, commonly used to seek relief from asthma cigarettes made of this mixtures are smoked.

The extract are prepared from roots and leaves , which are used as antifungal . Fresh leaves were washed through under tap water followed by sterilized water the leaves air , dried and were grinded with the help of pestle and mortar by taking (1:1 w/v) one gram of extract was added in 1ml distilled water separately for each plant extract and filtered through Muslin Cloth and 100% plant extract solution was prepared . The extracts were poured in the flasks plugged with cotton and heated at 100°C for 10 minutes to avoid contamination Karade V. M. and D. M. Sawant. (1999). The plants extracts Osuji, J.O. and Eunice, O.N. (2008). Different concentration (5,10,15,20%) of plant extracts was incorporated to potato dextrose medium agar for inoculation of the test pathogen in sterilized petridishes. The isolated pathogen was grown on potato dextrose agar medium was placed at the center of petridishes containing different concentration of the poisoned medium and incubated at 27±2°C for 6 days . Sharma R. R., Singh D. and Singh R.(2009). Radial growth (cm) of fungus was measured after inoculation till 6 days at an interval of 24 hrs.

Percentage of different leaf extract against fungi

Plant extracts	<i>Alternaria alternata</i>	<i>Aspergillus niger</i>	<i>Curvularia lunata</i>	<i>Phomopsis vexans</i>	<i>Fusarium solani</i>
<i>Azadirachtaindica</i>	62.15	74.25	64.20	49.30	55.57
<i>Argemone mexicana</i>	55.30	45.50	54.23	61.90	4.20
<i>Annona squamosa</i>	57.20	63.20	60.30	65.27	50.30
<i>Datura stramoniums</i>	28.20	30.52	55.40	22.15	23.15
<i>Calatropisprocera</i>	30.42	26.40	48.50	27.30	44.20
<i>Ipomoea fistulosa</i>	52.63	55.63	37.80	73.29	72.63
<i>Jatropacurcas</i>	36.15	46.15	30.12	46.20	35.12
<i>Lantana camara</i>	45.30	56.27	47.52	45.30	60.12
<i>Parthenium hysterothrus</i>	50.26	40.26	59.33	52.92	50.26
<i>Ocimumsantum</i>	60.00	3.20	25.15	16.20	60.00
<i>Nerium indicum</i>	32.15	29.15	42.50	40.15	20.30

RESULT AND DISCUSSION

During the study of present investigation the different plants have its own importance in the point view of antimicrobial compounds. For this investigation 11 plant extracts were taken to check the antifungal activity against the major Brinjal pathogenic fungi.

It is clear from the table that the highest disease efficacy was seen from *Datura* against the *Phomopsisvexans*. The lowest efficacy was seen in *Argemone mexicana* against the *Fusarium solani*. *Annona squamosa* shows the less efficacy against all selected five fungi. *Azadirachtaindica* shows more powerful against the *Curvularialunata*. *Ipomoea fistulosa* shows the lowest activity.

It is clear from the table different medicinal plants were selected for study during the investigation highest percent of a disease control efficacy (DCE) was observed in *Annona squamosa* leaf extract against the *Phomopsisvexans* i.e. (75.27%). The *Argemone mexicana* shows 64.20 % against the *Fusarium solani*. The *Azadirachtaindica* showing 64.20% efficacy against the *Curvularialunata* and 62.15% against the *Alternaria alternata*. Varshney vibha(2001). Among these 11 medicinal plants *Azadirachtaindica* showing highest 74.25% efficacy against the *Aspergillus niger*. The *Ipomoea fistulosa* also showing the notable efficiency against the *Phomopsisvexans* 73.29% and 72.63 % against the *Fusarium solani*. *Partheniumhysterothrus* shows disease control efficacy against the *Curvularialunata*. During the investigation lowest disease control efficacy was observed

References

1. Alemu Nega (2014) .Review on concepts in biological control of plant pathogens. *Journalof Biology, Agriculture and Healthcare*. 4, No. 27.
2. Hossain K.S. and Bashar M.A. (2011). *In vitro* effect of plant extracts, fungicides and antibiotics on the fungal isolates associated with damping-off disease of crucifers. *Journal of Agro forestry and Environment*. 5(2): 17-20.
3. Karade V. M. and D. M. Sawant. (1999).Effect of some plant extracts on the spore germination of *Alternaria alternata*. *Plant DiseaseResearch*, 14(1): 75.
4. Osuji, J.O. and Eunice, O.N. (2008).Evaluation of plant extracts for antifungal activity against *Sclerotium rolfsii* causing cocoyam cormel rot in storage. *Research Journal Agriculture Biological Science* 4(6): 784-787.
5. Sharma R. R., Singh D. and Singh R. (2009).Biological control of postharvest diseases of fruits and vegetables by microbial antagonists: a review. *Biological Control* 50, 205–221.
6. Varshney vibha(2001).Effect of plant extracts on *Drechsleragraminea*, the casual agent of simple of barley. *Indian Phytopathology*. 54. 32:378-381
7. Hossain KS and Bashar MA (2011). *In vitro* effect of plant extracts, fungicides and antibiotics on the fungal isolates associated with damping-off disease of crucifers. *Journal ofAgro forestry and Environment*. 5(2): 17-20.
8. Iqbal S. M., Ghafoor A., Ahmed Z. and Haqqani A. M. (2003). Pathogenecityand Fungicidal efficacy for *Sclerotinia* rot of Brinjal. *International Journal of Agriculture and Biology*. 5(4): 618-620.
9. Jagannathan R. and Narasimhan (1988).Effect of plant extracts on 2 fungal pathogen of Finger millet. *Indian Journal of Mycology and Plant Pathology*. 18(3): 250-254.