



Research Article

A STUDY OF (b, μ)-CONTINUITY, SLIGHTLY (b, μ)-CONTINUITY AND ITS RELATIONSHIPS

Santhi P¹ and Vijaya S²

¹Department of Mathematics, NPR College of Engineering and Technology, Natham, Dindigul District, Tamil Nadu, India

²Department of Mathematics, Sethu Institute of Technology, Kariyapatti, Virudhunagar District, Tamil Nadu, India

ARTICLE INFO

Article History:

Received 25th December, 2017

Received in revised form 18th

January, 2018 Accepted 14th February, 2018

Published online 28th March, 2018

ABSTRACT

The purpose of this paper is to study the concept related to λ-b-open sets, λ-b-closed sets and (b, μ)-continuity. Also some basic properties of slightly (b, μ)-continuous functions and connectedness and covering properties of slightly (b, μ)-continuous functions are investigated.

2010 Mathematics Subject Classification: 57C10, 57C08, 57C05

Key words:

(b, μ)-continuity, slightly (b, μ)-continuity, (λ-LC, μ)-continuity.

Copyright©2018 Santhi P and Vijaya S. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

In [2, 3, 4], Császár founded the theory of generalized topological spaces, and studied the extremely elementary character of these classes. Especially he introduced the notions of continuous functions on generalized topological spaces, and investigated characterizations of generalized continuous functions (= (λ, μ)-continuous functions in [3]). In [6, 7, 8], Min introduced the notions of weak (λ, μ)-continuity, almost (λ, μ)-continuity, (α, μ)-continuity, (σ, μ)-continuity, (π, μ)-continuity and (β, μ)-continuity on generalized topological spaces.

In this paper, we study the concept related to λ-b-open sets, λ-b-closed sets and (b, μ)-continuity. Also some basic properties of slightly (b, μ)-continuous functions and connectedness and covering properties of slightly (b, μ)-continuous functions are investigated.

2.Preliminaries

Definition

Let X be a nonempty set and λ be a collection of subsets of X. Then λ is called a generalized topology (briefly GT) on X if φ ∈ λ and G_i ∈ λ for i ∈ I ≠ φ implies G = ∪_{i ∈ I} G_i ∈ λ.

We say λ is strong if X ∈ λ, and we call the pair (X, λ) a generalized topological space (briefly GTS) on X.

The elements of λ are called λ-open sets and their complements are called λ-closed sets. The generalized closure of a subset S of X, denoted by c_λ(S), is the intersection of λ-closed sets including S. And the interior of S, denoted by i_λ(S), is the union of λ-open sets contained in S.

Definition 2.2

Let (X, λ) be a generalized topological space and A ⊂ X. Then A is said to be

1. λ-semi-open [3] if A ⊂ c_λ(i_λ(A)),
2. λ-preopen [3] if A ⊂ i_λ(c_λ(A)),
3. λ-α-open [3] if A ⊂ i_λ(c_λ(i_λ(A))),
4. λ-β-open [3] if A ⊂ c_λ(i_λ(c_λ(A))),
5. λ-b-open [12] if A ⊆ c_λ(i_λ(A)) ∪ i_λ(c_λ(A))

The complement of λ-semi-open (resp. λ-preopen, λ-α-open, λ-β-open, λ-b-open) is said to be λ-semi-closed (resp. λ-preclosed, λ-α-closed, λ-β-closed, λ-b-closed).

Let us denote by σ(λ_X) (briefly σ_X or σ) the class of all λ-semi-open sets on X, by π(λ_X) (briefly π_X or π) the class of all λ-preopen sets on X, by α(λ_X) (briefly α_X or

*Corresponding author: Santhi P

Department of Mathematics, NPR College of Engineering and Technology, Natham, Dindigul District, Tamil Nadu, India

α) the class of all λ - α -open sets on X , by $\beta(\lambda_X)$ (briefly β_X or β) the class of all λ - β -open sets on X , by $b(\lambda_X)$ (briefly b_X or b) the class of all λ - b -open sets on X . Obviously [9] $\lambda_X \subset \alpha(\lambda_X) \subset \sigma(\lambda_X) \subset \beta(\lambda_X)$ and $\lambda_X \subset \alpha(\lambda_X) \subset \pi(\lambda_X) \subset \beta(\lambda_X)$.

Lemma 2.3 [3]

Let (X, λ) be a generalized topological space and $A \subset X$. Then A is λ - α -open if and only if it is λ -semi-open and λ -preopen.

Definition 2.4 [10]

A GTS (X, λ) is said to be λ -extremally disconnected if the λ -closure of every λ -open set of X is λ -open in X .

Definition 2.5 [10]

Let (X, λ) be a generalized topological space and $\mathcal{B} \subset X$. Let $\phi \in \mathcal{B}$. Then \mathcal{B} is called a Base for λ if $\{\cup \mathcal{B}' : \mathcal{B}' \subset \mathcal{B}\} = \lambda$. We also say that λ is generated by \mathcal{B} .


Definition 2.6

Let (X, λ) and (Y, μ) be GTS's. Then a function $f : (X, \lambda) \rightarrow (Y, \mu)$ is said to be

1. (λ, μ) -continuous [4] if for each μ -open set U in Y , $f^{-1}(U)$ is λ -open in X ,
2. (α, μ) -continuous [7] if for each μ -open set U in Y , $f^{-1}(U)$ is λ - α -open in X ,
3. (σ, μ) -continuous [7] if for each μ -open set U in Y , $f^{-1}(U)$ is λ -semi-open in X ,
4. (π, μ) -continuous [7] if for each μ -open set U in Y , $f^{-1}(U)$ is λ -preopen in X ,
5. (β, μ) -continuous [7] if for each μ -open set U in Y , $f^{-1}(U)$ is λ - β -open in X .

Remark 2.7 [7]

Let f be a function between GTS's (X, λ) and (Y, μ) . Then we have the following implications.



Remark 2.8 [3]

Let (X, λ) be a generalized topological space and $A \subset X$. The λ -closure (resp. λ - α -closure, λ -semi-closure, λ -pre-closure, λ - β -closure, λ - b -closure) of a subset A of X , denoted by $c_\lambda(A)$ (resp. $c_\alpha(A)$, $c_\sigma(A)$, $c_\pi(A)$, $c_\beta(A)$, $c_b(A)$), is the intersection of all λ -closed (resp. λ - α -closed, λ -semi-closed, λ -pre-closed, λ - β -closed, λ - b -closed) sets containing A .

The λ -interior (resp. λ - α -interior, λ -semi-interior, λ -preinterior, λ - β -interior, λ - b -interior) of a subset A of X , denoted by $i_\lambda(A)$ (resp. $i_\alpha(A)$, $i_\sigma(A)$, $i_\pi(A)$, $i_\beta(A)$, $i_b(A)$), is the union of all λ -open (resp. λ - α -open, λ -semi-open, λ -preopen, λ - β -open, λ - b -open) sets contained in A .

Lemma 2.9 [4]

For a subset A of a GTS (X, λ) , we have

1. $X - i_\lambda(X - A) = c_\lambda(A)$,
2. $X - c_\lambda(X - A) = i_\lambda(A)$.

Lemma 2.10 [4]

Let A be a subset in a GTS (X, λ) . Then

1. $i_\lambda(c_\lambda(i_\lambda(c_\lambda(A)))) = i_\lambda(c_\lambda(A))$ and $c_\lambda(i_\lambda(c_\lambda(i_\lambda(A)))) = c_\lambda(i_\lambda(A))$.
2. $(i_\lambda(c_\lambda(A)))' = c_\lambda(i_\lambda(A'))$ and $(c_\lambda(i_\lambda(A)))' = i_\lambda(c_\lambda(A'))$.

Theorem 2.11 [3]

For any subset A in a GTS (X, λ) , the following statements are true.

1. $c_\sigma(A) = A \cup i_\lambda(c_\lambda(A))$ and $i_\sigma(A) = A \cap c_\lambda(i_\lambda(A))$.
2. $c_\pi(A) \supseteq A \cup c_\lambda(i_\lambda(A))$ and $i_\pi(A) \subseteq A \cap i_\lambda(c_\lambda(A))$.

Lemma 2.12 [4]

Let $f : (X, \lambda) \rightarrow (Y, \mu)$ be a function. For sets A and B of (X, λ) and (Y, μ) respectively, the following statements hold:

1. $f(f^{-1}(B)) \subseteq B$;
2. $f^{-1}(f(A)) \supseteq A$;
3. $f(A') \supseteq (f(A))'$;
4. $f^{-1}(B') = (f^{-1}(B))'$;
5. if f is injective, then $f^{-1}(f(A)) = A$;
6. if f is surjective, then $f(f^{-1}(B)) = B$;
7. if f is bijective, then $f(A') = (f(A))'$.

Definition 2.13 [5]

A set μ in a GTS (X, λ) is called λ -clopen if it is both λ -open and λ -closed.

Definition 2.14 [5]

A subset A in a GTS (X, λ) is said to be λ -locally λ -closed set (briefly λ -LC set) if $A = M \cap N$, where M is a λ -open set and N is a λ -closed set.

Definition 2.15 [11]

A GTS (X, λ) is said to be λ -connected if it cannot be expressed as the union of two nonempty, disjoint λ -open sets.

Definition 2.16 [1]

Let $f : (X, \lambda) \rightarrow (Y, \mu)$ be a function from a GTS (X, λ) to a GTS (Y, μ) . Then the function $g : X \rightarrow X \times Y$ defined by $g(x) = (x, f(x))$ is called the λ -graph function of f . Recall that for a function $f : (X, \lambda) \rightarrow (Y, \mu)$, the subset $\{(x, f(x)) : x \in X\} \subseteq X \times Y$ is called the λ -graph of f and is denoted by $G(f)$.

Definition 2.17 [13]

Let (X, λ) and (Y, μ) be GTS's. Then a function $f : (X, \lambda) \rightarrow (Y, \mu)$ is said to be almost (α, μ) -continuous if for each μ -regular open set U in Y , $f^{-1}(U)$ is λ - α -open in X .

3. λ - b -closed sets and (b, μ) -continuity

In this section, we obtain some new results which are related to λ - b -open sets, λ - b -closed sets and (b, μ) -continuity.

Definition 3.1

Let (X, λ) and (Y, μ) be GTS's. Then a function $f : (X, \lambda) \rightarrow (Y, \mu)$ is said to be

1. (b, μ) -continuous if for each μ -open set U in Y , $f^{-1}(U)$ is λ -b-open in X ,
2. $(\lambda$ -LC, $\mu)$ -continuous if for each μ -open set U in Y , $f^{-1}(U)$ is λ -locally λ -closed in X .

Remark3.2

Any union of λ -b-open sets is a λ -b-open set.

Theorem3.3

For a GTS (X, λ) , the following properties are equivalent.

1. (X, λ) is λ -extremally disconnected.
2. $i_\lambda(A)$ is λ -closed for every λ -closed subset A of X .
3. $c_\lambda(i_\lambda(A)) \subseteq i_\lambda(c_\lambda(A))$ for every subset A of X .
4. Every λ -semi-open set is λ -preopen.
5. The λ -closure of every λ - β -open subset of X is λ -open.
6. Every λ - β -open set is λ -preopen.
7. For every subset A of X , A is λ - α -open if and only if it is λ -semi-open.

Proof

1. \Rightarrow (ii): Let $A \subseteq X$ be a λ -closed set. Then A' is λ -open in X . By (i), $c_\lambda(A') = (i_\lambda(A))'$ is λ -open. Thus, $i_\lambda(A)$ is λ -closed.
2. \Rightarrow (iii): Let A be any set of X . Then $i_\lambda(A')$ is λ -closed in X . By (ii), $i_\lambda(i_\lambda(A'))$ is λ -closed in X . Therefore $i_\lambda(c_\lambda(A')) = c_\lambda(i_\lambda(A'))'$ is λ -closed in X and $c_\lambda(i_\lambda(A))$ is λ -open in X and hence $c_\lambda(i_\lambda(A)) \subseteq i_\lambda(c_\lambda(A))$.
3. \Rightarrow (iv): Let A be λ -semi-open. By (iii), we have $A \subseteq c_\lambda(i_\lambda(A)) \subseteq i_\lambda(c_\lambda(A))$. Thus A is λ -preopen.
4. \Rightarrow (v): Let A be a λ - β -open set. Then $c_\lambda(A)$ is λ -semi-open. By (iv), $c_\lambda(A)$ is λ -preopen. Thus, $c_\lambda(A) \subseteq i_\lambda(c_\lambda(A))$ and hence $c_\lambda(A)$ is λ -open.
5. \Rightarrow (vi): Let A be λ - β -open set. By (v), $c_\lambda(A) = i_\lambda(c_\lambda(A))$. Thus, $A \subseteq c_\lambda(A) = i_\lambda(c_\lambda(A))$ and hence A is λ -preopen.
6. \Rightarrow (vii): Let A be a λ -semi-open set. Since a λ -semi-open set is λ - β -open, then by (vi), it is λ -preopen. Since A is λ -semi-open and λ -preopen, A is λ - α -open.
7. \Rightarrow (i): Let A be a λ -open set of X . Then $c_\lambda(A)$ is λ -semi-open and by (vii) $c_\lambda(A)$ is λ - α -open. Therefore $c_\lambda(A) \subseteq i_\lambda(c_\lambda(i_\lambda(c_\lambda(A)))) = i_\lambda(c_\lambda(A))$ and hence $c_\lambda(A) = i_\lambda(c_\lambda(A))$. Hence $c_\lambda(A)$ is λ -open and (X, λ) is λ -extremally disconnected.

Theorem3.4

For a subset A of a GTS (X, λ) , then the following are equivalent.

1. A is a λ -locally λ -closed set.
2. $A = M \cap c_\lambda(A)$ for some λ -open set M .

Proof

- a. \Rightarrow (ii): Since A is a λ -locally λ -closed set. Then $A = M \cap N$ where M is λ -open and N is λ -closed. So $A \subseteq M$ and $A \subseteq N$. Hence $c_\lambda(A) \subseteq c_\lambda(N)$. Therefore $A \subseteq M \cap c_\lambda(A) \subseteq M \cap c_\lambda(N) = M \cap N = A$. Thus $A = M \cap c_\lambda(N)$.

- b. \Rightarrow (i): Since $A = M \cap c_\lambda(A)$ for some λ -open set M . Let $c_\lambda(A) = N$ be a λ -closed. We obtain $A = M \cap N$ where M is λ -open and N is λ -closed. Hence A is a λ -locally λ -closed set.

Theorem3.5

Let (X, λ) be a λ -extremally disconnected space and $A \subseteq X$, the following properties are equivalent.

1. A is a λ -open set.
2. A is λ - α -open and a λ -locally λ -closed set.
3. A is λ -preopen and a λ -locally λ -closed set.
4. A is λ -semi-open and a λ -locally λ -closed set.
5. A is λ -b-open and a λ -locally λ -closed set.

Proof

- a. \Rightarrow (ii): It follows from the fact that every λ -open set is λ - α -open and λ -locally λ -closed set.
- b. \Rightarrow (iii), (ii) \Rightarrow (iv), (iii) \Rightarrow (v) and (iv) \Rightarrow (v): Obvious.
- c. \Rightarrow (i): Suppose that A is a λ -b-open set and a λ -locally λ -closed set. It follows that $A \subseteq c_\lambda(i_\lambda(A)) \cup i_\lambda(c_\lambda(A))$. Since A is a λ -locally λ -closed set, then there exists a λ -open set M such that $A = M \cap c_\lambda(A)$. It follows that $A \subseteq M \cap [c_\lambda(i_\lambda(A)) \cup i_\lambda(c_\lambda(A))] = [M \cap c_\lambda(i_\lambda(A))] \cup [M \cap i_\lambda(c_\lambda(A))] \subseteq [M \cap i_\lambda(c_\lambda(A))] \cup [M \cap i_\lambda(c_\lambda(A))] = i_\lambda(M \cap c_\lambda(A)) \cup i_\lambda(M \cap c_\lambda(A)) = i_\lambda(A) \cup i_\lambda(A) = i_\lambda(A)$. Thus, $A \subseteq i_\lambda(A)$ and hence A is a λ -open set in X .

Theorem3.6

For a subset A of a GTS (X, λ) , if $A = c_\pi(A) \cap c_\sigma(A)$, then A is a λ -b-closed set.

Proof

Let $A = c_\pi(A) \cap c_\sigma(A) \supseteq (A \cup c_\lambda(i_\lambda(A))) \cap (A \cup i_\lambda(c_\lambda(A))) \supseteq c_\lambda(i_\lambda(A)) \cap i_\lambda(c_\lambda(A))$. This implies $c_\lambda(i_\lambda(A)) \cap i_\lambda(c_\lambda(A)) \subseteq A$. Thus A is λ -b-closed set in X .

Theorem

Let $f: (X, \lambda) \rightarrow (Y, \mu)$ where (X, λ) is a λ -extremally disconnected space and (Y, μ) is a GTS. Then the following properties are equivalent.

- a. f is a (λ, μ) -continuous.
- b. f is (α, μ) -continuous and a $(\lambda$ -LC, $\mu)$ -continuous.
- c. f is (π, μ) -continuous and a $(\lambda$ -LC, $\mu)$ -continuous.
- d. f is (σ, μ) -continuous and a $(\lambda$ -LC, $\mu)$ -continuous.
- e. f is (b, μ) -continuous and a $(\lambda$ -LC, $\mu)$ -continuous.

Proof

It is obvious from Theorem 3.5.

4.Slightly (b, μ) -Continuous Functions

In this section, basic properties of slightly (b, μ) -continuous functions and connectedness and covering properties of slightly (b, μ) -continuous functions are investigated.

Definition 4.1

Let (X, λ) and (Y, μ) be generalized topological spaces. A function $f : (X, \lambda) \rightarrow (Y, \mu)$ is said to be slightly (b, μ) -continuous if for each point $x \in X$ and each μ -clopen set M in Y containing $f(x)$, there exists a λ -b-open set N in X containing x such that $f(N) \subseteq M$.

Theorem

For a function $f : (X, \lambda) \rightarrow (Y, \mu)$, the following statements are equivalent:

- a. f is slightly (b, μ) -continuous;
- b. for every μ -clopen set A in Y , $f^{-1}(A)$ is λ -b-open;
- c. for every μ -clopen set A in Y , $f^{-1}(A)$ is λ -b-closed;
- d. for every μ -clopen set A in Y , $f^{-1}(A)$ is λ -b-clopen.

Proof

- a. \Rightarrow (2): Let A be a μ -clopen set in Y and let $x \in f^{-1}(A)$. Since $f(x) \in A$, by (1), there exists a λ -b-open set M in X containing x such that $M \subseteq f^{-1}(A)$. We obtain that $f^{-1}(A) = \cup M (x \in f^{-1}(A))$. Thus, $f^{-1}(A)$ is λ -b-open.
- b. \Rightarrow (3): Let A be a λ -clopen set in Y . Then, A' is λ -clopen. By (2), $f^{-1}(A') = (f^{-1}(A))'$ is λ -b-open. Thus, $f^{-1}(A)$ is λ -b-closed.
- c. \Rightarrow (4): It can be shown easily.
- a. \Rightarrow (1): Let A be a λ -clopen set in Y containing $f(x)$. By (4), $f^{-1}(A)$ is λ -b-clopen. Take $M = f^{-1}(A)$. Then, $f(M) \subseteq A$. Hence, f is slightly (b, μ) -continuous.

Remark 4.3

Obviously (b, μ) -continuity implies slightly (b, μ) -continuity. The following example shows that this implication is not reversible.

Example 4.4

Let $X = \{a, b\}$, $Y = \{p, q\}$ and λ, μ are sets defined as follows: $\lambda = \{\phi, X, a\}$ and $\mu = \{\phi, Y, q\}$. Then the function $f : (X, \lambda) \rightarrow (Y, \mu)$ defined by $f(a) = p$, $f(b) = q$ is slightly (b, μ) -continuous but not (b, μ) -continuous.

Theorem 4.5

Suppose that Y has a base consisting of λ -clopen sets. If $f : (X, \lambda) \rightarrow (Y, \mu)$ is slightly (b, μ) -continuous, then f is (b, μ) -continuous.

Proof

Let $x \in X$ and let A be a μ -open set in Y containing $f(x)$. Since Y has a base consisting of λ -clopen sets, there exists a λ -clopen set M containing $f(x)$ such that $M \subseteq A$. Since f is slightly (b, μ) -continuous, then there exists a λ -b-open set N in X containing x such that $f(N) \subseteq M \subseteq A$. Thus, f is (b, μ) -continuous.

Theorem 4.6

Let $f : (X, \lambda) \rightarrow (Y, \mu)$ be a function and let $g : X \rightarrow X \times Y$ be the λ -graph function of f , defined by $g(x) = (x, f(x))$ for every $x \in X$. If g is slightly (b, μ) -continuous, then f is slightly (b, μ) -continuous.

Proof

Let M be μ -clopen set in Y , then $X \times M$ is μ -clopen set in $X \times Y$. Since g is slightly (b, μ) -continuous, then $f^{-1}(M) = g^{-1}(X \times M)$ is λ -b-open in X . Thus, f is slightly (b, μ) -continuous.

Definition 4.7

A filter base Λ is said to be λ -b-convergent to a point x in X if for any λ -b-open set M in X containing x , there exists a set $N \in \Lambda$ such that $N \subseteq M$.

Definition 4.8

A filter base Λ is said to be λ -co-convergent to a point x in X if for any λ -clopen set M in X containing x , there exists a set $N \in \Lambda$ such that $N \subseteq M$.

Theorem 4.9

If a function $f : (X, \lambda) \rightarrow (Y, \mu)$ is slightly (b, μ) -continuous, then for each point $x \in X$ and each filter base Λ in X λ -b-converging to x , the filter base $f(\Lambda)$ is λ -co-convergent to $f(x)$.

Proof

Let $x \in X$ and Λ be any filter base in X λ -b-converging to x . Since f is slightly (b, μ) -continuous, then for any μ -clopen set M in Y containing $f(x)$, there exists a λ -b-open set N in X containing x such that $f(N) \subseteq M$. Since Λ is λ -b-converging to x , there exists a $P \in \Lambda$ such that $P \subseteq N$. This means that $f(P) \subseteq M$ and therefore the filter base $f(\Lambda)$ is λ -co-convergent to $f(x)$.

Definition 4.10

A GTS (X, λ) is said to be λ -b-connected if it cannot be expressed as the union of two nonempty, disjoint λ -b-open sets.

Theorem 4.11

If $f : (X, \lambda) \rightarrow (Y, \mu)$ is slightly (b, μ) -continuous surjective function and X is λ -b-connected space, then Y is μ -connected space.

Proof

Suppose that Y is not μ -connected space. Then there exists nonempty disjoint μ -open sets M and N such that $Y = M \cup N$. Therefore, M and N are μ -clopen sets in Y . Since f is slightly (b, μ) -continuous, then $f^{-1}(M)$ and $f^{-1}(N)$ are λ -b-closed and λ -b-open in X . Moreover, $f^{-1}(M)$ and $f^{-1}(N)$ are nonempty disjoint and $X = f^{-1}(M) \cup f^{-1}(N)$. This shows that X is not λ -b-connected. This is a contradiction. By contradiction, Y is μ -connected.

Definition 4.12

A GTS (X, λ) is said to be

1. λ -b-compact if every λ -b-open cover of X has a finite subcover.
2. countably λ -b-compact if every λ -b-open countably cover of X has a finite subcover.
3. λ -b-Lindelof if every cover of X by λ -b-open sets has a countable subcover.
4. λ -mildly compact if every λ -clopen cover of X has a finite subcover.
5. λ -mildly countably compact if every λ -clopen countably cover of X has a finite subcover.

6. λ -mildly Lindelof if every cover of X by λ -clopen sets has a countable subcover.

Theorem4.13

Let $f: (X, \lambda) \rightarrow (Y, \mu)$ be a slightly (b, μ) -continuous surjection. Then the following statements hold:

- if X is λ -b-compact, then Y is λ -mildly compact.
- if X is λ -b-Lindelof, then Y is λ -mildly Lindelof.
- if X is countably λ -b-compact, then Y is λ -mildly countably compact.

Proof

Let $\{A_\alpha : \alpha \in I\}$ be any μ -clopen cover of Y . Since f is slightly (b, μ) -continuous, then $\{f^{-1}(A_\alpha) : \alpha \in I\}$ is a λ -b-open cover of X . Since X is λ -b-compact, there exists a finite subset I_0 of I such that $X = \cup\{f^{-1}(A_\alpha) : \alpha \in I_0\}$. Thus, we have $Y = \cup\{A_\alpha : \alpha \in I_0\}$ and Y is λ -mildly compact. The other proofs are similar.

Definition4.14

A GTS (X, λ) is said to be

- λ -b-closed-compact if every λ -b-closed cover of X has a finite subcover.
- countably λ -b-closed-compact if every countable cover of X by λ -b-closed sets has a finite subcover.
- λ -b-closed-Lindelof if every cover of X by λ -b-closed sets has a countable subcover.

Theorem4.15

Let $f: (X, \lambda) \rightarrow (Y, \mu)$ be a slightly (b, μ) -continuous surjection. Then the following statements hold:

- if X is λ -b-closed-compact, then Y is λ -mildly compact.
- if X is λ -b-closed-Lindelof, then Y is λ -mildly Lindelof.
- if X is countably λ -b-closed-compact, then Y is λ -mildly countably compact.

Proof

It can be obtained similarly as the previous theorem.

5.The Relationships

In this section, we investigate the relationships between slightly (b, μ) -continuous functions and separation axioms and the relationships between slightly (b, μ) -continuity and λ -graphs.

Definition 5.1

A GTS (X, λ) is said to be λ -b- T_1 if for each pair of distinct points x and y of X , there exist λ -b-open sets M and N containing x and y respectively such that $y \notin M$ and $x \notin N$.

Definition5.2

A GTS (X, λ) is said to be λ -co- T_1 if for each pair of distinct points x and y of X , there exist λ -clopen sets M and N containing x and y respectively such that $y \notin M$ and $x \notin N$.

Definition5.3

If $f: (X, \lambda) \rightarrow (Y, \mu)$ is a slightly (b, μ) -continuous injection and Y is μ -co- T_1 , then X is λ -b- T_1 .

Proof

Suppose that Y is μ -co- T_1 . For any distinct points x and y in X , there exist λ -clopen sets M, N in Y such that $f(x) \in M, f(y) \notin M, f(x) \notin N$ and $f(y) \in N$. Since f is slightly (b, μ) -continuous,

$f^{-1}(M)$ and $f^{-1}(N)$ are λ -b-open sets in X such that $x \in f^{-1}(M), y \notin f^{-1}(M), x \notin f^{-1}(N)$ and $y \in f^{-1}(N)$. This shows that X is λ -b- T_1 .

Definition5.4

A GTS (X, λ) is said to be λ -b- T_2 (λ -b-Hausdorff) if for each pair of distinct points x and y in X , there exist disjoint λ -b-open sets M and N in X such that $x \in M$ and $y \in N$.

Definition5.5

A GTS (X, λ) is said to be λ -co- T_2 (λ -co-Hausdorff) if for each pair of distinct points x and y in X , there exist disjoint λ -clopen sets M and N in X such that $x \in M$ and $y \in N$.

Theorem5.6

If $f: (X, \lambda) \rightarrow (Y, \mu)$ is a slightly (b, μ) -continuous injection and Y is μ -co- T_2 , then X is λ -b- T_2 .

Proof

For any pair of distinct points x and y in X , there exist disjoint μ -clopen sets M and N in Y such that $f(x) \in M$ and $f(y) \in N$. Since f is slightly (b, μ) -continuous, $f^{-1}(M)$ and $f^{-1}(N)$ is λ -b-open in X containing x and y respectively. We have $f^{-1}(M) \cap f^{-1}(N) = \emptyset$. This shows that X is λ -b- T_2 .

Definition5.7

A GTS (X, λ) is called λ -co-regular (respectively strongly λ -b-regular) if for each λ -clopen (respectively λ -b-closed) set A and each point $x \notin A$, there exist disjoint λ -open sets M and N such that $A \subseteq M$ and $x \in N$.

Definition5.8

A GTS (X, λ) is said to be λ -co-normal (respectively strongly λ -b-normal) if for every pair of disjoint λ -clopen (respectively λ -b-closed) sets A and B in X , there exist disjoint λ -open sets M and N such that $A \subseteq M$ and $B \subseteq N$.

Theorem5.9

If f is slightly (b, μ) -continuous injective λ -open function from a strongly λ -b-regular space X onto a GTS (Y, μ) then Y is μ -co-regular.

Proof

Let A be μ -clopen set in Y and $y \notin A$. Take $y = f(x)$. Since f is slightly (b, μ) -continuous, by Theorem 4.3(3) $f^{-1}(A)$ is a λ -b-closed set. Take $B = f^{-1}(A)$. We have $x \notin B$. Since X is strongly λ -b-regular, there exist disjoint λ -open sets M and N such that $B \subseteq M$ and $x \in N$. We obtain that $A = f(B) \subseteq f(M)$ and $y = f(x) \in f(N)$ such that $f(M)$ and $f(N)$ are disjoint μ -open sets. This shows that Y is μ -co-regular.

Theorem5.10

If f is slightly (b, μ) -continuous injective λ -open function from a strongly λ -b-normal space (X, λ) onto a GTS (Y, μ) then Y is μ -co-normal.

Proof

Let A and B be disjoint μ -clopen sets in Y . Since f is slightly (b, μ) -continuous, $f^{-1}(A)$ and $f^{-1}(B)$ are λ -b-closed sets. Take $M = f^{-1}(A)$ and $N = f^{-1}(B)$. We have $M \cap N = \emptyset$. Since X is strongly λ -b-normal, there exist disjoint λ -open sets P and Q such that $M \subseteq P$ and $N \subseteq Q$. We obtain that $A = f(M) \subseteq$

$f(P)$ and $B = f(N) \subseteq f(Q)$ such that $f(P)$ and $f(Q)$ are disjoint μ -open sets. Thus, Y is λ -co-normal.

Definition5.11

A λ -graph $G(f)$ of a function $f : (X, \lambda) \rightarrow (Y, \mu)$ is said to be λ -b-co-closed if for each $(x, y) \in (X \times Y) \setminus G(f)$, there exist a λ -b-open set M in X containing x and a μ -clopen set N in Y containing y such that $(M \times N) \cap G(f) = \emptyset$.

Lemma 5.12

A λ -graph $G(f)$ of a function $f : (X, \lambda) \rightarrow (Y, \mu)$ is λ -b-co-closed in $X \times Y$ if and only if for each $(x, y) \in (X \times Y) \setminus G(f)$, there exist a λ -b-open set M in X containing x and a μ -clopen set N in Y containing y such that $f(M) \cap N = \emptyset$.

Theorem5.13

If $f : (X, \lambda) \rightarrow (Y, \mu)$ is slightly (b, μ) -continuous and Y is μ -co-Hausdorff, then $G(f)$ is λ -b-co-closed in $X \times Y$.

Proof

Let $(x, y) \in (X \times Y) \setminus G(f)$, then $f(x) \neq y$. Since Y is μ -co-Hausdorff, there exist μ -clopen sets M and N in Y with $f(x) \in M$ and $y \in N$ such that $M \cap N = \emptyset$. Since f is slightly (b, μ) -continuous, there exists a λ -b-open set P in X containing x such that $f(P) \subseteq M$. Therefore, we obtain $y \in N$ and $f(P) \cap N = \emptyset$. This shows that $G(f)$ is λ -b-co-closed.

Theorem5.14

If $f : (X, \lambda) \rightarrow (Y, \mu)$ is (b, μ) -continuous and Y is μ -co- T_1 , then $G(f)$ is λ -b-co-closed in $X \times Y$.

Proof

Let $(x, y) \in (X \times Y) \setminus G(f)$, then $f(x) \neq y$. Since Y is μ -co- T_1 , there exist μ -clopen sets M and N in Y with $f(x) \in M$ and $y \in N$ such that there exists a λ -clopen set N' in Y such that $f(x) \notin N'$ and $y \in N'$. Since f is (b, μ) -continuous, there exists a λ -b-open set $f^{-1}(A)$ in X containing x such that $f(M) \subseteq N'$. Therefore, we obtain that $f(M) \cap (N') = \emptyset$ and N' is μ -clopen set containing y . This shows that $G(f)$ is λ -b-co-closed in $X \times Y$.

Theorem5.15

Let $f : (X, \lambda) \rightarrow (Y, \mu)$ has a λ -b-co-closed graph $G(f)$. If f is injective, then X is λ -b- T_1 .

Proof

Let x and y be any two distinct points of X . Then, we have $(x, f(y)) \in (X \times Y) \setminus G(f)$. By definition of λ -b-co-closed graph, there exist a λ -b-open set M in X and a λ -clopen set N in Y such that $x \in M$, $f(y) \in N$ and $f(M) \cap N = \emptyset$; hence $M \cap f^{-1}(N) = \emptyset$. Therefore, we have $y \notin M$. This implies that X is λ -b- T_1 .

Definition5.16

A function $f : (X, \lambda) \rightarrow (Y, \mu)$ is called always λ -b-open if the image of each λ -b-open set in X is μ -b-open set in Y .

Theorem5.17

Let $f : (X, \lambda) \rightarrow (Y, \mu)$ has a λ -b-co-closed graph $G(f)$. If f is surjective always λ -b-open function, then Y is μ -b- T_2 .

Proof

Let y and z be any distinct points of Y . Since f is surjective, $f(x) = y$ for some $x \in X$ and $(x, z) \in (X \times Y) \setminus G(f)$. By λ -b-co-closedness of graph $G(f)$, there exists a λ -b-open set M in X and a μ -clopen set N in Y such that $x \in M$, $z \in N$ and $(M \times N) \cap G(f) = \emptyset$. Then, we have $f(M) \cap N = \emptyset$. Since f is always λ -b-open, then $f(M)$ is μ -b-open such that $f(x) = y \in f(M)$. This implies that Y is μ -b- T_2 .

References

1. S.-Z. Bai and Y.-P Zuo, On g - \square -irresolute functions, *Acta Math. Hungar.*, 130 (4) (2011), 382-389., doi: 10.1007/s10474-010-0014-x.
2. A. Császár, Generalized open sets, *Acta Math. Hungar.*, 75 (1-2) (1997), 65-87, doi: 10.1023/A:1006582718102.
3. A. Császár, Generalized open sets in generalized topologies, *Acta Math. Hungar.*, 106(1-2) (2005), 53-66, doi: 10.1007/s10474-005-0005-5.
4. A. Császár, Generalized topology, generalized continuity, *Acta Math. Hungar.*, 96 (4) (2002), 351-357, doi: 10.1023/A:1019713018007.
5. R. Jamunarani and P.Jeyanthi, Regular sets in generalized topological spaces, *Acta Math. Hungar.*, 135 (4) (2012), 342-349, doi: 10.1007/s10474-011-0179-y.
6. W. K. Min, Almost continuity on generalized topological spaces, *Acta Math. Hungar.*, 125 (1-2) (2009), 121-125, doi: 10.1007/s10474-009-8230-y.
7. W. K. Min, Generalized continuous functions defined by generalized open sets on generalized topological spaces, *Acta Math. Hungar.*, 128 (4) (2009), 299-306, doi: 10.1007/s10474-009-9037-6.
8. W. K. Min, Some results on generalized topological spaces and generalized systems, *Acta Math. Hungar.*, 108 (1-2) (2005), 171-181, doi: 10.1007/s10474-005-0218-7.
9. W. K. Min, Weak continuity on generalized topological spaces, *Acta Math. Hungar.*, 124 (1-2) (2008), 73-81, doi: 10.1007/s10474-008-8152-0.
10. R. D. Sarma, On extremally disconnected generalized topologies, *Annales Univ. Budapest, Sectio Math*, 47 (2004), 91-96, 151-161.
11. R. X. Shen, A note on generalized connectedness, *Acta Math. Hungar.*, 122(3) (2008), 231-235, doi: 10.1007/s10474-008-8009-6.
12. P. Sivagami, Remarks on γ -interior, *Acta Math. Hungar.*, 119 (1-2) (2008), 81-94, doi: 10.1007/s1047-007-7007-4.
13. S. Vijaya and P.Santhi, Characterization of almost (\square, μ) -continuous functions and its properties, *International Science and Technology Journal*, Volume 7, Issue 3, (2018), 67-74, doi:16.10089. ISTJ.2018. V713. 285311.2610
